



Houtwal 101 Postal Code 5509
KK Veldhoven
HOLLAND

Doxin-50-WSP

Powder for oral administration

Composition:

Contains per gram powder:

Doxycycline hyclate.....500 mg.
Excipients ad1 g.

Description:

Doxycycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria like Bordetella, Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella, Staphylococcus and Streptococcus spp. Doxycycline is also active against Chlamydia, Mycoplasma and Rickettsia spp. The action of doxycycline is based on inhibition of bacterial protein synthesis. Doxycycline has a great affinity to the lungs and is therefore especially useful for treatment of bacterial respiratory infections.

Indications:

Gastrointestinal and respiratory infections caused by doxycycline sensitive micro-organisms, like Bordetella, Campylobacter, Chlamydia, E. coli, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in calves, goats, poultry, sheep and swine.

Contra-indications:

Hypersensitivity to tetracyclines.

Administration to animals of a seriously impaired hepatic function.

Concurrent administration with penicillins, cephalosporins, quinolones and cycloserine. Administration to animals with an active microbial digestion.

Side effects:

Discoloration of teeth in young animals may occur. Hypersensitivity reactions may occur.

Dosage:

For oral administration:

Calves, goats and sheep : Twice daily 1 g per 100 kg body weight for 3 - 5 days.

Poultry and swine : 100 g per 500 - 1000 litres of drinking water for 3 - 5 days.

Note: for pre-ruminant calves, lambs and kids only.

Withdrawal times:

- For meat:

Calves, goats and sheep : 14 days

Swine : 8 days

Poultry : 7 days

Packaging:

Sachet of 100 g and jar of 1000 g.