



Houtwal 101 Postal Code 5509
KK Veldhoven
HOLLAND

FLORXIN[®] 10

Composition per ml:

Florfenicol 100 mg .
Excipients: N-methyl-2-pyrrolidone, propylene glycol, PEG ad1 ml .

Pharmaceutical form:

Oral solution for administration via drinking water, consisting of a clear, yellowish solution.

Withdrawal period:

For poultry meat: 2 day from the last administration.

Not permitted for use in laying bird producing eggs for human consumption.

Indication:

In poultry: for treatment of respiratory and digestive bacterial infections produced by susceptible bacteria: Pasteurella multocida, Bordetella Bronchiseptica, Salmonella spp., Escherichia coli, Haemophilus Paragalinarum, Staphylococcus spp., streptococcus spp., Ornithobacterium rhinotracheale and Corynebacterium pyogenes.

Contraindications:

Do not administer in animals with known hypersensitivity at florfenicol.

Dosage, Route (s) and method of administration:

The product is administered via drinking water in poultry as following:

In poultry: the product is administered via drinking water in dosage of 20 mg florfenicol/kg b.w.: 0.2 ml of product/kg b.w./day in birds under 4 weeks of age, and 0.4 ml of product/kg b.w./day in birds over 4 weeks of age, for 3-5 days.