



Houtwal 101 Postal Code 5509
KK Veldhoven
HOLLAND

N-Flox20

Solution for oral administration

Composition

Contains per ml:

Norfloxacin.....200 mg.
Excipients ad 1 ml.

Description:

Norfloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like *Campylobacter*, *E. coli*, *Haemophilus*, *Pasteurella*, *Salmonella*, and *Mycoplasma* spp.

Indications:

Gastrointestinal, respiratory and urinary tract infections caused by norfloxacin sensitive micro-organisms, like *Campylobacter*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella* and *Salmonella* spp. in calves, goats, poultry, sheep and swine.

Contra-indications:

Hypersensitivity to norfloxacin.

Administration to animals with a seriously impaired hepatic and/or renal function. Concurrent administration of tetracyclines, chloramphenicol, macrolides and lincosamides.

Side effects:

Hypersensitivity reactions.

Administration to juvenile animals can lead to arthropathy.

Dosage:

For oral administration:

Calves, goats and sheep : Twice daily 10 ml per 75 - 150 kg body weight for 3 - 5 days.

Poultry : 1 litre per 1500 - 4000 litres of drinking water for 3 - 5 days.

Swine : 1 litre per 1000 - 3000 litres of drinking water for 3 - 5 days.

Note: for pre-ruminant calves, lambs and kids only.

Withdrawal times:

- For meat

Calves, goats, sheep and swine : 8 days.

Poultry : 12 days.

Packaging:

Bottle of 100, 500 and 1000 ml.